

Boundary non-crossings of Brownian pillow

Enkelejd Hashorva ¹

University of Bern

Abstract. Let $B_0(s, t)$ be a Brownian pillow with continuous sample paths, and let $h, u : [0, 1]^2 \rightarrow \mathbb{R}$ be two measurable functions. In this paper we derive upper and lower bounds for the boundary non-crossing probability $\psi(u; h) := \mathbf{P}\{B_0(s, t) + h(s, t) \leq u(s, t), \forall s, t \in [0, 1]\}$. Further we investigate the asymptotic behaviour of $\psi(u; \gamma h)$ with γ tending to ∞ , and solve a related minimisation problem.

Key words and phrases: Boundary non-crossing probability; Brownian pillow with trend; large deviations; smallest concave majorant; Reproducing Kernel Hilbert Space; small ball probabilities.

1 Introduction

Let $B_0(s, t), s, t \in [0, 1]$ be a Brownian pillow with continuous sample paths. Its covariance function K is a product of two covariance functions defined by

$$K((s_1, t_1), (s_2, t_2)) = K_1(s_1, t_1)K_2(s_2, t_2), \quad s_i, t_i \in [0, 1], i = 1, 2,$$

with $K_i(s, t) = \min(s, t) - ts$, $i = 1, 2$ the covariance function of a Brownian bridge.

Our concern in this article is the boundary non-crossing probability

$$\psi(u; h) := \mathbf{P}\{B_0(s, t) + h(s, t) \leq u(s, t), \forall s, t \in [0, 1]\}, \quad (1.1)$$

with h a trend function and u a measurable boundary function.

When considering Brownian bridge and Brownian motion the corresponding non-crossing probability can be explicitly calculated if h and u are polygonal lines, see e.g., Goovaerts and Teunen (1993), Wang and Pötzelberger (1997), Novikov et al. (1999), Janssen and Kunz (2004), Borovkov and Novikov (2005) and the references therein. Such explicit formulae are not available in our setup of the multi-parameter processes.

Our novel results presented below are:

a) upper and lower bounds for $\psi(u; h)$, b) a large deviation type result for the boundary non-crossing probability $\psi(u; \gamma h)$ with $\gamma \rightarrow \infty$, and c) we solve a related minimisation problem.

We comment briefly the result mentioned in b). Given a function $g : [0, \infty)^2 \rightarrow \mathbb{R}$ we denote by g'' its partial derivative obtained by differentiating both components, provided it exists. From the large deviation theory (see e.g., Lifshits (1995) or Ledoux (1996)) for any positive constant c and any trend function $h : [0, 1]^2 \rightarrow \mathbb{R}$ with a square integrable partial derivative h'' (i.e., $\int_{[0, 1]^2} (h''(s, t))^2 ds dt < \infty$) we obtain

$$\lim_{\gamma \rightarrow \infty} 2\gamma^{-2} \ln \mathbf{P}\left\{ \sup_{s, t \in [0, 1]} (B_0(s, t) + \gamma h(s, t)) \leq c \right\} = - \int_{[0, 1]^2} (\underline{h}''(s, t))^2 ds dt \in (-\infty, 0], \quad (1.2)$$

¹Department of Mathematical Statistics and Actuarial Science, University of Bern, Sidlerstrasse 5, CH-3012 Bern, Switzerland;enkelejd.hashorva@stat.unibe.ch

September 26, 2008

with $\underline{h}$ the solution of the minimisation problem

$$\inf_{g \geq \underline{h}} \int_{[0,1]^2} (g''(s,t))^2 ds dt, \quad (1.3)$$

where the functions $g : [0, 1]^2 \rightarrow \mathbb{R}$ in the minimisation problem are assumed to possess a square integrable partial derivative g'' , and g, h vanish on the boundary of $[0, 1]^2$.

Compared to (1.2) our new result is a sharper asymptotic estimate of the boundary non-crossing probability of interest. In the special case h being a product of two concave functions $h_1, h_2 : [0, 1] \rightarrow [0, \infty)$ with $h_i(0) = h_i(1) = 0, i = 1, 2$ we show (see below (4.5))

$$\begin{aligned} & \mathbf{P} \left\{ \sup_{s,t \in [0,1]} (B_0(s,t) + \gamma h_1(s) h_2(t)) \leq c \right\} \\ &= \exp \left(-\frac{\gamma^2}{2} \prod_{i=1,2} \int_{[0,1]} (h'_i(x))^2 \lambda(dx) + c\gamma \prod_{i=1,2} [h'_i(1) - h'_i(0)] + z(\gamma) \right), \end{aligned} \quad (1.4)$$

where

$$-A\gamma^{2/3} \ln^3 \gamma \leq z(\gamma) \leq \ln \mathbf{P} \left\{ \sup_{s,t \in [0,1]} B_0(s,t) \leq c \right\}$$

holds for all large γ with a positive constant A not depending on γ . Here h'_i is a right continuous version of the derivative of $h_i, i = 1, 2$ and λ is the Lebesgue measure on $[0, 1]$.

We derive (1.4) utilising a known small ball result for Brownian pillow. Indeed the small ball problem for both the Brownian pillow and the Brownian sheet is investigated by several authors, see Kuelbs and Li (1992), Talagrand (1994), Csáki et al. (2000), Khoshnevisan and Pemantle (2000), Li and Shao (2001), Koning and Protasov (2003), Fill and Torcaso (2004), Gao et al. (2004), Gao and Li (2006, 2007) or Karol' et al. (2008) among many other references.

A consequence of the Gaussian shift inequality (see Li and Kuelbs (1998)) and (1.4) is the following bound (set D for the set of all concave functions $f : [0, 1] \rightarrow [0, \infty)$)

$$\mathbf{P} \left\{ \sup_{s,t \in [0,1]} B_0(s,t) \leq c \right\} \leq \inf_{h \in D} \Phi \left(c^2 \left(\frac{h'(1) - h'(0)}{\int_0^1 (h'(x))^2 \lambda(dx)} \right)^2 \right), \quad (1.5)$$

with Φ the distribution function of a Gaussian random variable with mean 0 and variance 1. Since the upper bound in (1.5) is not smaller than $1/2$, the above inequality is of some interest provided that $\psi(0; c) \in (1/2, 1)$.

Organisation of the paper: In the next section we present some notation and preliminary results. The main results are discussed in Section 3. Section 4 explains the simple situation where the trend function h is a product of two trend functions. Proofs of all the results are relegated to Section 5 followed by a short Appendix with two results on the Riemann-Stieltjes integral.

2 Preliminaries

We introduce first a Hilbert spaces related to the covariance function of the Brownian pillow, which can also be seen as tensor product of Hilbert spaces related to the covariance function of the Brownian bridge. Then we provide a result utilised in solving the minimisation problem (1.3).

The Reproducing Kernel Hilbert Space (RKHS) related to the covariance function of a Brownian pillow, denoted by $\mathcal{H}_2^0$, is given by

$$\mathcal{H}_2^0 := \left\{ h : [0, 1]^2 \rightarrow \mathbb{R} \mid \exists h'' \in L_2([0, 1]^2, \lambda^2), \text{ with } h(s, t) = \int_{[0, s] \times [0, t]} h''(x, y) \lambda^2(dx, dy), \right. \\ \left. h(0, s) = h(1, s) = h(t, 0) = h(t, 1) = 0, \quad \forall s, t \in [0, 1] \right\},$$

where $L_2([0, 1]^2, \lambda^2)$ is the set of all real functions defined on $[0, 1]^2$ which are square integrable with respect to the Lebesgue measure λ^2 on $[0, 1]^2$. The inner product is

$$\langle h_1, h_2 \rangle = \int_{[0, 1]^2} h_1''(x, y) h_2''(x, y) \lambda^2(dx, dy), \quad h_1, h_2 \in \mathcal{H}_2^0$$

and the corresponding norm of $h \in \mathcal{H}_2^0$ is $\|h\| := \langle h, h \rangle^{1/2}$.

As shown in Koning and Protasov (2003) another approach to deal with $\mathcal{H}_2^0$ is to construct this Hilbert space as the tensor product of two RKHS, i.e., $\mathcal{H}_2^0 = \mathcal{H}_1^0 \otimes \mathcal{H}_1^0$ with $\mathcal{H}_1^0$ the RKHS of the covariance function of the Brownian bridge defined by

$$\mathcal{H}_1^0 := \left\{ h : [0, 1] \rightarrow \mathbb{R} \mid \exists h' \in L_2([0, 1], \lambda), \text{ with } h(s) = \int_{[0, s]} h'(x) \lambda(dx), \quad h(0) = h(1) = 0 \right\},$$

where $L_2([0, 1], \lambda)$ is the set of all real functions defined on $[0, 1]$ that are square integrable with respect to λ . The inner product of $\mathcal{H}_1^0$ is

$$\langle h_1, h_2 \rangle = \int_{[0, 1]} h_1'(x) h_2'(x) \lambda(dx), \quad h_1, h_2 \in \mathcal{H}_1^0$$

and the corresponding norm is denoted again by $\|\cdot\|$. Any element $h \in \mathcal{H}_2^0$ can be identified by $h_1, h_2 \in \mathcal{H}_1^0$ so that $h = h_1 \otimes h_2$ (see Koning and Protasov (2003)).

In the following for any trend function $h \in \mathcal{H}_2^0$ we write by h'' its right continuous derivative.

Lemma 2 in Janssen and Hülya (2008) is crucial for our next result. Define the closed convex sets

$$V := \{h \in \mathcal{H}_2^0 : h(s, t) \leq 0, \forall s, t \in [0, 1]\}, \quad W := \{h \in \mathcal{H}_2^0 : h(s, t) \geq 0, \forall s, t \in [0, 1]\},$$

and let $\widetilde{V}, \widetilde{W}$ be the polar cones of V and W , respectively defined by

$$\widetilde{V} := \{h \in \mathcal{H}_2^0 : \langle h, v \rangle \leq 0, \forall v \in V\}, \quad \widetilde{W} := \{h \in \mathcal{H}_2^0 : \langle h, v \rangle \geq 0, \forall v \in W\}.$$

Further denote by $BV_H(T), T \subset \mathbb{R}^2$ the class of functions $f : T \rightarrow \mathbb{R}$ which have bounded variation in the sense of Hardy (see e.g., Adams and Clarkson (1934), Móricz (2002)).

Lemma 2.1 *Let $h \in \mathcal{H}_2^0$ be a given function and let $V_{p,h}, \widetilde{V}_{p,h}$ be the unique projections of h into V and the polar cone $\widetilde{V}$, respectively.*

a) If $\widetilde{V}_{p,h}''$ is a right continuous partial derivative of $\widetilde{V}_{p,h}$ such that $\widetilde{V}_{p,h}'' \in BV_H([0, 1]^2)$, then for any function $g : [0, 1]^2 \rightarrow [0, \infty)$ Riemann-Stieltjes integrable with respect to $\widetilde{V}_{p,h}''$ the Riemann-Stieltjes integral

$I(g) := \int_{[0,1]^2} g(s,t) d\tilde{V}_{p,h}''(s,t)$ satisfies $I(g) \geq 0$.

b) We have

$$h = V_{p,h} + \tilde{V}_{p,h}, \quad \langle V_{p,h}, \tilde{V}_{p,h} \rangle = 0. \quad (2.1)$$

c) If $h = h_1 + h_2$ with $h_1 \in V, h_2 \in \tilde{V}$ such that $\langle h_1, h_2 \rangle = 0$, then $h_1 = V_{p,h}$ and $h_2 = \tilde{V}_{p,h}$.

d) The unique solution $\underline{h}$ of the minimisation problem

$$\min_{g \geq h, g \in \mathcal{H}_2^0} \|g\| \quad (2.2)$$

is $\underline{h} = \tilde{V}_{p,h}$ satisfying further $\|\underline{h}\| = \min\{\|g\| : g \in \tilde{V}, g \geq h\}$.

We note in passing that a similar decomposition to (2.1) can be stated for $h \in \mathcal{H}_2^0$ in terms of the unique projections $W_{p,h}, \tilde{W}_{p,h}$ of h into W and the polar cone $\tilde{W}$, respectively. Furthermore, b), c) hold for some general Hilbert space.

We write alternatively $\underline{h}, \bar{h}$ instead of $\tilde{V}_{p,h}, \tilde{W}_{p,h}$. The above lemma immediately implies

$$\bar{h}(s,t) \leq h(s,t) \leq \underline{h}(s,t), \quad \forall s, t \in [0,1], \quad \text{and } \|h\| \geq \max(\|\underline{h}\|, \|\bar{h}\|), \quad \forall h \in \mathcal{H}_2^0. \quad (2.3)$$

Furthermore, for any two functions $h, q \in \mathcal{H}_2^0, q \geq h$ (1.3) and Lemma 2.1 yield

$$\|q\| \geq \|h\|, \quad (2.4)$$

provided that $\underline{h} = h, \underline{q} = q$.

3 Main Results

Let $B_0(s,t), s, t \in [0,1]$ be a Brownian pillow with continuous sample paths, and let $h \in \mathcal{H}_2^0$ be a given trend function. For some measurable boundary function $u : [0,1]^2 \rightarrow \mathbb{R}$ we define the boundary non-crossing probability $\psi(u; h)$ as in (1.1). Throughout the rest of the paper we assume that $\psi(u; 0) \in (0,1)$. Since $h \in \mathcal{H}_2^0$ the Cameron-Martin formula (see e.g., Kuelbs (1976), Lifshits (1995), Li and Kuelbs (1998) or Li and Shao (2001)) implies

$$\psi(u; h) = \exp\left(-\frac{1}{2}\|h\|^2\right) \mathbf{E}\left\{\exp\left(\int_{[0,1]^2} h''(s,t) dB_0(s,t)\right) \mathbf{1}(B_0(s,t) \leq u(s,t), \forall s, t \in [0,1])\right\}, \quad (3.1)$$

where $\mathbf{1}(\cdot)$ is the indicator function.

Li and Kuelbs (1998) show that the Cameron-Martin translation implies important shift inequalities for some general Gaussian processes. Applying their Theorem 1' we have

$$\Phi(\theta - \|h\|) \leq \psi(u; h) \leq \Phi(\theta + \|h\|), \quad (3.2)$$

where Φ is the Gaussian distribution function on $\mathbb{R}$ with mean 0 and variance 1, and θ is such that $\Phi(\theta) = \psi(u; 0)$. When $\|h\|$ is small the lower and the upper bounds in (3.2) are close to the non-crossing probability of interest, since $\lim_{\gamma \rightarrow 0} \psi(u; \gamma h) = \psi(u; 0) = \Phi(\theta)$. As $\gamma \rightarrow \infty$ the upper bound in (3.2)

tends to 1, whereas the lower bound and $\psi(u; \gamma h)$ tend to 0. Note in passing that as in Pötzelberger and Wang (2001) we obtain

$$|\psi(u; \gamma h) - \psi(u; 0)| \leq 2\Phi(\gamma\|h\|/2) - 1 \leq \frac{\gamma\|h\|}{\sqrt{2\pi}}, \quad \forall \gamma \in (0, \infty). \quad (3.3)$$

One important criteria which we will look at when discussing bounds for the non-crossing probability of interest is their performance for both small or large trend functions. In our first result below we provide upper and lower bounds for the boundary non-crossing probability $\psi(u; h)$. If we consider further the trend function γh , then the bounds perform well when $\gamma \rightarrow 0$.

Proposition 3.1 *Let $h, u : [0, 1]^2 \rightarrow \mathbb{R}$ be two measurable functions such that $\psi(u; 0) \in (0, 1)$. If $h \in \mathcal{H}_2^0$, then we have*

$$\Phi(\theta - \|\underline{h}\|) \leq \psi(u; h) \leq \Phi(\theta + \|\bar{h}\|), \quad \theta := \Phi^{-1}(\psi(u; 0)), \quad (3.4)$$

with $\underline{h}, \bar{h}$ as defined in Section 2 and Φ^{-1} the inverse of Φ . Furthermore

$$-\frac{\|\underline{h}\|}{\sqrt{2\pi}} \leq \psi(u; h) - \psi(u; 0) \leq \frac{\|\bar{h}\|}{\sqrt{2\pi}}. \quad (3.5)$$

When $h \neq \underline{h}$ or $h \neq \bar{h}$ in view of (2.3) we see that (3.5) yields better bounds than (3.3). By (3.5) we obtain

$$-\gamma \frac{\|\underline{h}\|}{\sqrt{2\pi}} \leq \psi(u; \gamma h) - \psi(u; 0) \leq \gamma \frac{\|\bar{h}\|}{\sqrt{2\pi}}, \quad \forall \gamma > 0, \quad (3.6)$$

which is of some interest when γ tends to 0 since both the lower and the upper bounds converge to 0. As mentioned in the Introduction if γ tends to infinity, then we have the logarithmic asymptotic behaviour

$$\lim_{\gamma \rightarrow \infty} 2\gamma^{-2} \ln \psi(u; \gamma h) = -\|\underline{h}\|^2, \quad \forall h \in \mathcal{H}_2^0, \quad (3.7)$$

with $\underline{h}$ the unique solution of the minimisation problem (2.2).

Next, we derive explicit upper and lower bounds for $\psi(u; h)$, which perform better asymptotically (for trend function becoming large) as those implied by (3.4).

Proposition 3.2 *Let $h \in \mathcal{H}_2^0$ be a given trend function, and let $u, l : [0, 1]^2 \rightarrow \mathbb{R}$ be two measurable functions. If the partial derivative $\underline{h}''$ of the projection of h into its polar cone satisfies $\underline{h}'' \in BV_H([0, 1]^2)$ and is right continuous, then*

$$\underline{h} := \inf_{g \geq h, g \in \tilde{V}, g \in BV_H([0, 1]^2)} g \quad (3.8)$$

and further $\underline{h}$ is the smallest majorant of h such that its right continuous partial derivative belongs to $BV_H([0, 1]^2)$ and generates a finite positive measure.

Moreover, if the Riemann-Stieltjes integral $\int_{[0, 1]^2} v(s, t) d\underline{h}''(s, t)$ is finite for both $v = l$ and $v = u$ and $\psi(u; 0) \in (0, 1)$, then

$$\psi(u; h) \leq \psi(u; h - \underline{h}) \exp\left(-\frac{1}{2}\|\underline{h}\|^2 + \int_{[0, 1]^2} u(s, t) d\underline{h}''(s, t)\right) \quad (3.9)$$

and

$$\begin{aligned} \psi(u; h) &\geq \mathbf{P}\{l(s, t) \leq B_0(s, t) \leq u(s, t), \forall s, t \in [0, 1]\} \\ &\quad \times \exp\left(-\frac{1}{2}\|\underline{h}\|^2 + \int_{[0,1]^2} l(s, t) d\underline{h}''(s, t)\right) \end{aligned} \quad (3.10)$$

are valid.

Remarks: a) If $u(s, t) := c \in (0, \infty), \forall s, t \in [0, 1]$, then (3.9) implies

$$\psi(c; h) \leq \psi(c; h - \underline{h}) \exp\left(-\frac{1}{2}\|\underline{h}\|^2 + c[\underline{h}''(1, 1) - \underline{h}''(1, 0) - \underline{h}''(0, 1) + \underline{h}''(0, 0)]\right). \quad (3.11)$$

A lower bound for $\psi(c; h)$ is derived utilising (3.10) with $l(s, t) := -c, \forall s, t \in [0, 1]$.

b) As in the proof of Proposition 3.2 it can be shown that if the trend function $h \in \mathcal{H}_2^0$ is such that its right continuous partial derivative h'' satisfies $h'' \in BV_H([0, 1]^2)$ and furthermore h'' generates a positive measure on $[0, 1]^2$, then the unique solution of the minimisation problem (2.2) is $\underline{h} = h$.

c) An upper bound for $\psi(u; h)$ is the discrete boundary non-crossing probability

$$\psi_n(u; h) := \mathbf{P}\{B_0(s_i, t_i) + h(s_i, t_i) \leq u(s_i, t_i), \forall (s_i, t_i) \in T_n\},$$

with $T_n := \{(s_i, t_i), i = 1, \dots, n\} \subset [0, 1]^2$. Hashorva (2005b) shows the asymptotic behaviour (considering the Brownian bridge) of the corresponding discrete boundary non-crossing probability.

Next, we discuss the asymptotic behaviour of $\psi(u; \gamma h)$ when $\gamma \rightarrow \infty$. Exact asymptotics of the non-crossing probabilities of the Brownian motion with trend is derived in Hashorva (2005a) which was motivated by a large deviation type result obtained in Bischoff et al. (2003). As in Bischoff et al. (2005) we expect that our novel asymptotic result will have some implications for statistical applications.

Proposition 3.3 Let $h, \underline{h}, u$ be as in Proposition 3.2. Suppose that there exist functions $u_\varepsilon \in \mathcal{H}_{2,\varepsilon}^0, \varepsilon > 0$ such that $\|u_\varepsilon\| = O(1/\varepsilon)$ and

$$\lim_{\varepsilon \rightarrow 0} u_\varepsilon(s, t) = u(s, t), \quad u_\varepsilon(s, t) \leq u(s, t) - \varepsilon, \quad \forall s, t \in [0, 1]. \quad (3.12)$$

If the Riemann-Stieltjes integral $I_\varepsilon := \int_{[0,1]^2} u_\varepsilon(s, t) d\underline{h}''(s, t)$ exists and $|I_\varepsilon| \leq M \in (0, \infty), \forall \varepsilon > 0$, then

$$\lim_{\varepsilon \rightarrow 0} I_\varepsilon = I := \int_{[0,1]^2} u(s, t) d\underline{h}''(s, t), \quad |I| \leq M, \quad (3.13)$$

and

$$\psi(u; \gamma h) = \exp\left(-\frac{\gamma^2}{2}\|\underline{h}\|^2 + \gamma \int_{[0,1]^2} u(s, t) d\underline{h}''(s, t) + z(\gamma)\right) \quad (3.14)$$

holds, where for all large γ

$$-A\gamma^{2/3} \ln^3 \gamma \leq z(\gamma) \leq \ln \mathbf{P}\{B_0(s, t) \leq u(s, t), \forall s, t \in [0, 1] : \underline{h}(s, t) = h(s, t)\}, \quad (3.15)$$

with a positive constant A not depending on γ .

In view of the above asymptotics and (3.4) we obtain a simple upper bound for $\psi(u; 0)$.

Corollary 3.4 *Let $u : [0, 1]^2 \rightarrow \mathbb{R}$ be a measurable function satisfying the assumptions of Proposition 3.3. Then we have*

$$\psi(u; 0) \leq \inf_{h \in \mathcal{H}_2^0, h'' \in BV_H([0, 1]^2) : \|\underline{h}\| > 0} \Phi \left(\|\underline{h}\|^{-1} \int_{[0, 1]^2} u(s, t) d\underline{h}''(s, t) \right). \quad (3.16)$$

Remarks: a) If the function u in Proposition 3.3 satisfies $u(s, t) > \mu \in (0, \infty), \forall s, t \in [0, 1]$ where (s, t) belongs to the boundary of $[0, 1]^2$, and there exist functions $w_\varepsilon : [0, 1]^2 \rightarrow \mathbb{R}, \varepsilon > 0$ such that $uw_\varepsilon \in \mathcal{H}_2^0, \varepsilon > 0$, then we may define u_ε in Proposition 3.3 by $u_\varepsilon := uw_\varepsilon - \varepsilon, \varepsilon > 0$. When u is a positive constant, then functions $u_\varepsilon, \varepsilon > 0$ satisfying the assumption of Proposition 3.3 can be easily constructed. If u_ε is continuous, then the Riemann-Stieltjes integral $I_\varepsilon := \int_{[0, 1]^2} u_\varepsilon(s, t) d\underline{h}''(s, t)$ in Proposition 3.3 is finite.

b) When $\underline{h}''$ is almost surely continuous with respect to the Lebesgue measure λ^2 , then instead of assuming that $\underline{h}$ has a bounded variation in the sense of Hardy (Lemma 2.1, Proposition 3.2 and 3.3) we may impose the weaker assumption $\underline{h}$ has a bounded variation in the sense of Vitali (see Appendix below and Lemma 6.2).

c) Our results can be easily extended to the d -dimensional setup considering $B_0(s_1, \dots, s_d), s_i \in [0, 1], i \leq d$ a Brownian pillow with continuous sample paths. The term $\ln^3 \gamma$ in (3.15) should then be replaced by $\ln^{2d-1} \gamma$.

d) Similar results can be stated for considering instead of B_0 a Brownian sheet $B(s, t), s, t \in [0, \infty)$ with continuous sample paths. For instance Proposition 3.2 holds with $\underline{h}$ the solution of the minimisation problem (1.3) where g, h have square integrable partial derivatives satisfying further $g(0, s) = h(0, s) = h(t, 0) = g(t, 0) = 0, s, t \in [0, \infty)$.

4 Product Trend Functions

As demonstrated in the previous section the non-crossing probability $\psi(u; h)$ can be bounded by some functions which depend on the solution of the minimisation problem (2.2). We discuss below an instance where the solution of (2.2) can be easily determined. Let therefore $h_1, h_2 \in \mathcal{H}_1^0$, and let $B_0(s), s \in [0, 1]$ denote a Brownian bridge with continuous sample paths. If $u_1, u_2 : [0, 1] \rightarrow \mathbb{R}$ are two measurable functions with $u_i(0), u_i(1) > 0, i = 1, 2$, then we have (see Bischoff and Hashorva (2005))

$$\begin{aligned} & \mathbf{P}\{B_0(s) + h_i(s) \leq u_i(s), \forall s \in [0, 1]\} \\ & \leq \mathbf{P}\{B_0(s) \leq u_i(s) + \tilde{h}_i(s) - h_i(s), \forall s \in [0, 1]\} \exp\left(-\frac{1}{2}\|\tilde{h}_i\|^2 + \int_{[0, 1]} u_i(s) d(-\tilde{h}_i'(s))\right), \end{aligned}$$

where $\tilde{h}_i, i = 1, 2$ is the smallest concave majorant of h_i and $\tilde{h}_i'$ is a right continuous derivative of $\tilde{h}_i$. Furthermore, $\tilde{h}_i$ is the unique solution of the minimisation problem

$$\min_{g \in \mathcal{H}_1^0, g \geq h_i} \|g\|, \quad i = 1, 2. \quad (4.1)$$

Set in the following $h(s, t) := h_1(s)h_2(t)$, $\tilde{h}(s, t) := \tilde{h}_1(s)\tilde{h}_2(t)$, $s, t \in [0, 1]$, and write $h = h_1 \times h_2$, $\tilde{h} = \tilde{h}_1 \times \tilde{h}_2$. In the next lemma we show that for special trend functions the unique solution of (2.2) with $h = h_1 \times h_2 \in \mathcal{H}_2^0$ is simply $\tilde{h}$.

Lemma 4.1 *Let $h := h_1 \times h_2$, $h_1, h_2 \in \mathcal{H}_1^0$, and denote by $\tilde{h}_i, i = 1, 2$ the smallest concave majorant of $h_i, i = 1, 2$. If*

$$\tilde{h}(s, t) \geq h(s, t), \quad \forall s, t \in [0, 1], \quad (4.2)$$

then the unique solution $\underline{h}$ of (1.3) is $\underline{h} := \tilde{h}$.

Clearly, (4.2) holds if h_1, h_2 are both non-negative functions. In the special case that also u is a product function we have the following immediate result.

Corollary 4.2 *Let $h_i, \tilde{h}_i, i = 1, 2$ satisfy the assumption of Lemma 4.1, and let $u_i, l_i : [0, 1] \rightarrow \mathbb{R}, i = 1, 2$ be measurable functions. If the Riemann-Stieltjes integral $\int_{[0,1]} v_i(s) d(-\tilde{h}'_i(s))$ is a finite constant for $i = 1, 2$ and $v_i = l_i$ or $v_i = u_i$, then we have*

$$\psi(u; h) \leq \psi(u; h - \tilde{h}) \exp\left(-\frac{1}{2}\|\tilde{h}_1\|^2\|\tilde{h}_2\|^2 + \prod_{i=1,2} \int_{[0,1]} u_i(s) d(-\tilde{h}'_i(s))\right), \quad (4.3)$$

with $h := h_1 \times h_2$, $\tilde{h} := \tilde{h}_1 \times \tilde{h}_2$, $u := u_1 \times u_2$, and further

$$\begin{aligned} \psi(u; h) &\geq \mathbf{P}\{l_1(s)l_2(t) \leq B(s, t) \leq u_1(s)u_2(t), \forall s, t \in [0, 1]\} \\ &\times \exp\left(-\frac{1}{2}\|\tilde{h}_1\|^2\|\tilde{h}_2\|^2 + \prod_{i=1,2} \int_{[0,1]} l_i(s) d(-\tilde{h}'_i(s))\right). \end{aligned} \quad (4.4)$$

Corollary 4.3 *Under the assumptions and the notation of Corollary 4.2 if further $\min_{s \in [0,1]} u_i(s) > C \in (0, \infty), i = 1, 2$ and $u_i, i = 1, 2$ are absolute continuous with u'_i satisfying $\int_{[0,1]} (u'_i(s))^2 \lambda(ds) < \infty$, then we have*

$$\psi(u_1 \times u_2; \gamma h_1 \times h_2) = \exp\left(-\frac{\gamma^2}{2}\|\tilde{h}_1\|^2\|\tilde{h}_2\|^2 + \gamma \prod_{i=1}^2 \int_{[0,1]} u_i(s) d(-\tilde{h}'_i(s)) + z(\gamma)\right) \quad (4.5)$$

with $z(\gamma)$ satisfying

$$-A\gamma^{2/3} \ln^3 \gamma \leq z(\gamma) \leq \ln \mathbf{P}\{B_0(s, t) \leq u_1(s)u_2(t), \forall s, t \in [0, 1] : \tilde{h}_1(s)\tilde{h}_2(t) = h_1(s)h_2(t)\}$$

for all large γ where A is a positive constant not depending on γ . Furthermore

$$\psi(u; 0) \leq \inf_{h_1, h_2 \in \mathcal{H}_1^0 : \|\tilde{h}_1\| \|\tilde{h}_2\| > 0} \Phi\left((\|\tilde{h}_1\| \|\tilde{h}_2\|)^{-1} \prod_{i=1,2} \int_{[0,1]} u_i(s) d(-\tilde{h}'_i(s))\right). \quad (4.6)$$

5 Proofs

Proof of Lemma 2.1: Let $g, h \in \mathcal{H}_2^0$ be two given functions. If $h'' \in BV_H([0, 1]^2)$ with h'' a right continuous partial derivative of h , then we have by (6.2) and the integration by parts formula (see Lemma 2 and Lemma 3 in Móricz (2002) and (6.1))

$$\langle g, h \rangle = \int_{[0,1]^2} g''(s, t) h''(s, t) \lambda^2(ds, dt)$$

$$\begin{aligned}
&= \int_{[0,1]^2} h''(s, t) dg(s, t) \\
&= \int_{[0,1]^2} g(s, t) dh''(s, t).
\end{aligned} \tag{5.1}$$

Consequently, for any $g \in V$ by the assumption on $\tilde{V}_{p,h}''$ we have $\langle g, \tilde{V}_{p,h} \rangle \leq 0$. Hence for any function $g : [0, 1]^2 \rightarrow [0, \infty)$ which is Riemann-Stieltjes integrable with respect to $\tilde{V}_{p,h}''$ on $[0, 1]^2$ we have for the corresponding Riemann-Stieltjes integral

$$\int_{[0,1]^2} g(s, t) d\tilde{V}_{p,h}'' \geq 0. \tag{5.2}$$

The proof of statements *b*) and *c*) follows immediately by Lemma 2 in Janssen and Hülya (2008).

We show next statement *d*). Let $\tilde{h} \in \mathcal{H}_2^0$ be a given function such that $\tilde{h} := g + h$ with $g(s, t) \geq 0, \forall s, t \in [0, 1]$. By the properties of $\tilde{V}_{p,h}$ we have $\langle \tilde{V}_{p,h}, g \rangle \geq 0$, hence we may write

$$\begin{aligned}
\|\tilde{h}\|^2 &= \|g + h\|^2 \\
&= \|\tilde{V}_{p,h} + g + h - \tilde{V}_{p,h}\|^2 \\
&= \|\tilde{V}_{p,h}\|^2 + 2\langle \tilde{V}_{p,h}, g + h - \tilde{V}_{p,h} \rangle + \|g + h - \tilde{V}_{p,h}\|^2 \\
&= \|\tilde{V}_{p,h}\|^2 + 2\langle \tilde{V}_{p,h}, g \rangle + 2\langle \tilde{V}_{p,h}, \tilde{V}_{p,h} \rangle + \|g + h - \tilde{V}_{p,h}\|^2 \\
&= \|\tilde{V}_{p,h}\|^2 + 2\langle \tilde{V}_{p,h}, g \rangle + \|g + h - \tilde{V}_{p,h}\|^2 \\
&\geq \|\tilde{V}_{p,h}\|^2 + 2\langle \tilde{V}_{p,h}, g \rangle \\
&\geq \|\tilde{V}_{p,h}\|^2.
\end{aligned}$$

Since further $\tilde{V}_{p,h}(s, t) \geq h(s, t), \forall s, t \in [0, 1]$ it follows that the solution of the minimisation problem (2.2) is $\tilde{V}_{p,h}$. Clearly, its solution is unique, thus the result follows. $\square$

Proof of Proposition 3.1: By (2.3) and (3.2) we see that (3.4) follows easily. The proof of (3.5) can be established along the lines of the proof of Lemma 5 in Janssen and Hülya (2008), thus the result. $\square$

Proof of Proposition 3.2: Let $V, \tilde{V}$ be as in Section 2, and let $\tilde{V}_{p,h}$ be the projection of h into the polar cone $\tilde{V}$. In view of statement *b*) of Lemma 2.1

$$h = V_{p,h} + \tilde{V}_{p,h}, \quad \|h\|^2 = \|\tilde{V}_{p,h}\|^2 + \|V_{p,h}\|^2.$$

Furthermore, $\psi(u; h) \geq \psi(u; \tilde{V}_{p,h})$. Next, applying the Cameron-Martin formula we obtain (set $\mathbf{1}_u(B_0(s, t)) := \mathbf{1}(B_0(s, t) \leq u(s, t), \forall s, t \in [0, 1])$)

$$\begin{aligned}
\psi(u; h) &= \exp\left(-\frac{1}{2}\|h\|^2\right) \mathbf{E}\left\{\exp\left(\int_{[0,1]^2} h''(s, t) dB_0(s, t)\right) \mathbf{1}_u(B_0(s, t))\right\} \\
&= \exp\left(-\frac{1}{2}\|\tilde{V}_{p,h}\|^2\right) \mathbf{E}\left\{\exp\left(-\frac{1}{2}\|V_{p,h}\|^2 + \int_{[0,1]^2} V_{p,h}''(s, t) dB_0(s, t)\right.\right. \\
&\quad \left.\left.+ \int_{[0,1]^2} \tilde{V}_{p,h}''(s, t) dB_0(s, t)\right) \mathbf{1}_u(B_0(s, t))\right\}.
\end{aligned}$$

Since $\tilde{V}_{p,h}'' \in BV_H([0,1]^2)$ is right continuous and $B_0(s,t)$ has continuous sample paths by the integration by parts formula (6.1) for the Riemann-Stieltjes integral we have almost surely

$$\int_{[0,1]^2} B_0(s,t) d\tilde{V}_{p,h}''(s,t) = \int_{[0,1]^2} \tilde{V}_{p,h}''(s,t) dB_0(s,t).$$

Consequently, we may further write (recall (5.2))

$$\begin{aligned} \psi(u; h) &= \mathbf{E} \left\{ \exp \left(-\frac{1}{2} \|V_{p,h}\|^2 + \int_{[0,1]^2} V_{p,h}''(s,t) dB_0(s,t) + \int_{[0,1]^2} B_0(s,t) d\tilde{V}_{p,h}''(s,t) \right) \mathbf{1}_u(B_0(s,t)) \right\} \\ &\leq \exp \left(-\frac{1}{2} \|\tilde{V}_{p,h}\|^2 + \int_{[0,1]^2} u(s,t) d\tilde{V}_{p,h}''(s,t) \right) \\ &\quad \mathbf{E} \left\{ \exp \left(-\frac{1}{2} \|V_{p,h}\|^2 + \int_{[0,1]^2} V_{p,h}''(s,t) dB_0(s,t) \right) \mathbf{1}_u(B_0(s,t)) \right\} \\ &= \exp \left(-\frac{1}{2} \|\tilde{V}_{p,h}\|^2 + \int_{[0,1]^2} u(s,t) d\tilde{V}_{p,h}''(s,t) \right) \psi(u; V_{p,h}). \end{aligned}$$

Clearly, by the definition $\psi(u; h) \geq \psi(u; \tilde{V}_{p,h})$. Applying (3.7) to $\psi(u; \gamma \tilde{V}_{p,h})$, $\gamma > 0$ we find

$$\ln \psi(u; \gamma h) = -(1 + o(1)) \frac{\gamma^2}{2} \|\tilde{V}_{p,h}\|^2, \quad \gamma \rightarrow \infty,$$

hence by (3.7) the unique solution of (2.2) equals $\tilde{V}_{p,h}$. Since $\tilde{V}_{p,h} \geq h$ and $\tilde{V}_{p,h} \in \tilde{V}$, then $\underline{h} = \tilde{V}_{p,h}$ and (3.8) follows.

We show next the last claim (3.10). Utilising again the Cameron-Martin formula we have

$$\begin{aligned} \psi(u; h) &\geq \psi(u; \underline{h}) \\ &\geq \mathbf{P} \{ l(s,t) \leq B_0(s,t) + \underline{h}(s,t) \leq u(s,t), \forall s, t \in [0,1] \} \\ &= \exp \left(-\frac{1}{2} \|\underline{h}\|^2 \right) \mathbf{E} \left\{ \exp \left(\int_{[0,1]^2} \underline{h}''(s,t) dB_0(s,t) \right) \mathbf{1} \{ l(s,t) \leq B_0(s,t) \leq u(s,t), \forall s, t \in [0,1] \} \right\} \\ &= \mathbf{P} \{ l(s,t) \leq B_0(s,t) \leq u(s,t), \forall s, t \in [0,1] \} \exp \left(-\frac{1}{2} \|\underline{h}\|^2 + \int_{[0,1]^2} l(s,t) d\underline{h}''(s,t) \right), \end{aligned}$$

hence the proof is established. $\square$

Proof of Proposition 3.3: Set next

$$\underline{h}_\epsilon(s,t) := \underline{h}(s,t) - u_\epsilon(s,t), \quad \forall s, t \in [0,1].$$

Applying Cameron-Martin formula we obtain

$$\begin{aligned} \psi(u; h) &\geq \psi(u; \underline{h}) \\ &= \mathbf{P} \{ B_0(s,t) + \underline{h}(s,t) \leq u(s,t), \forall s, t \in [0,1] \} \\ &\geq \mathbf{P} \{ B_0(s,t) + \underline{h}(s,t) \leq u_\epsilon(s,t) + \epsilon, \forall s, t \in [0,1] \} \\ &> \exp \left(-\frac{1}{2} \|\underline{h}_\epsilon\|^2 \right) \mathbf{E} \left\{ \exp \left(\int_{[0,1]^2} \underline{h}_\epsilon''(s,t) dB_0(s,t) \right) \mathbf{1} \{ -\epsilon \leq B_0(s,t) \leq \epsilon, \forall s, t \in [0,1] \} \right\}. \end{aligned}$$

Define the Gaussian random variable

$$Z := \int_{[0,1]^2} \underline{h}_\epsilon''(s,t) dB_0(s,t).$$

Clearly, Z has mean 0 and variance $\|\underline{h}_\epsilon\|^2$. For $\epsilon > 0$ small enough we have $\|\underline{h}_\epsilon\| \in (0, \infty)$. For any constant $C \in \mathbb{R}$ and ϵ small enough we may write

$$\begin{aligned}
& \mathbf{E}\{\exp(Z)\mathbf{1}(-\epsilon \leq B_0(s, t) \leq \epsilon, \forall s, t \in [0, 1])\} \\
&= \mathbf{E}\{\exp(Z)\mathbf{1}(-\epsilon \leq B_0(s, t) \leq \epsilon, \forall s, t \in [0, 1])[\mathbf{1}(Z < C) + \mathbf{1}(Z \geq C)]\} \\
&\geq \mathbf{E}\{\exp(Z)\mathbf{1}(-\epsilon \leq B_0(s, t) \leq \epsilon, \forall s, t \in [0, 1])\mathbf{1}(Z \geq C)\} \\
&\geq \exp(C)\mathbf{P}\{-\epsilon \leq B_0(s, t) \leq \epsilon, \forall s, t \in [0, 1], Z \geq C\} \\
&= \exp(C)\left[\mathbf{P}\left\{\sup_{s, t \in [0, 1]} |B_0(s, t)| < \epsilon\right\} - \mathbf{P}\{-\epsilon \leq B_0(s, t) \leq \epsilon, \forall s, t \in [0, 1], Z < C\}\right] \\
&\geq \exp(C)\left[\mathbf{P}\left\{\sup_{s, t \in [0, 1]} |B_0(s, t)| < \epsilon\right\} - \mathbf{P}\{Z \leq C\}\right] \\
&= \exp(C)[\mathbf{P}\left\{\sup_{s, t \in [0, 1]} |B_0(s, t)| < \epsilon\right\} - \Phi(C/\|\underline{h}_\epsilon\|)].
\end{aligned}$$

By the small ball asymptotic result (see Fill and Torcaso (2004), Gao and Li (2006, 2007), Karol' et al. (2008)) we have

$$\mathbf{P}\left\{\sup_{s, t \in [0, 1]} |B_0(s, t)| < \epsilon\right\} \geq \exp\left(-K \frac{\ln^3(1/\epsilon)}{\epsilon^2}\right)$$

for some positive constant K and all $\epsilon > 0$ small enough. Since

$$\|\underline{h}_\epsilon\|^2 = \|\underline{h}\|^2 - 2 \int_{[0, 1]^2} u_\epsilon(s, t) d\underline{h}''(s, t) + \|u_\epsilon\|^2 = O(1/\epsilon^2)$$

choosing $C := -K_* \|\underline{h}_\epsilon\| \ln^{3/2}(1/\epsilon)/\epsilon$, $K_* \in (0, \infty)$, $K_*^2 > K$ and using the Mills-Ratio asymptotics for Gaussian random variables for all $\epsilon > 0$ small enough and some positive constants c_1, c_2 we have

$$\mathbf{E}\{\exp(Z)\mathbf{1}(-\epsilon \leq B_0(s, t) \leq \epsilon, \forall s, t \in [0, 1])\} \geq \exp\left(-\frac{c_1}{\epsilon} - \frac{c_2 \ln^3(1/\epsilon)}{\epsilon^2}\right),$$

implying thus

$$\psi(u; h) \geq \exp\left(-\frac{1}{2}\|\underline{h}\|^2 + \int_{[0, 1]^2} u_\epsilon(s, t) d\underline{h}''(s, t) - \frac{c_1}{\epsilon} - \frac{c_2 \ln^3(1/\epsilon)}{\epsilon^2}\right).$$

Recalling that $\lim_{\epsilon \rightarrow 0} u_\epsilon(s, t) = u(s, t)$, $\forall s, t \in [0, 1]$ and $\|u_\epsilon\|^2 = O(1/\epsilon^2)$ we obtain utilising the result of Proposition 3.2 (set next $\epsilon := \gamma^{-1/3}$, $\gamma > 0$)

$$\psi(u; \gamma h) = \exp\left(-\frac{\gamma^2}{2}\|\underline{h}\|^2 + \gamma I + z(\gamma)\right), \quad \gamma \rightarrow \infty,$$

where $|I| \leq M$ with $I := \int_{[0, 1]^2} u(s, t) d\underline{h}''(s, t)$ and

$$-A\gamma^{2/3} \ln^3 \gamma \leq z(\gamma) \leq \ln \mathbf{P}\left\{B_0(s, t) \leq u(s, t), \forall s, t \in [0, 1] : \underline{h}(s, t) = h(s, t)\right\}$$

is satisfied for all γ large and A a positive constant not depending on γ . Hence the result follows. $\square$

Proof of Lemma 4.1: Set $V := \{h \in \mathcal{H}_2^0 : h(s, t) \leq 0, \forall s, t \in [0, 1]\}$ and $\underline{h} := \tilde{h}_1 \times \tilde{h}_2$. By the assumptions the function $g := \underline{h} - h_1 \times h_2$ belongs to V . Furthermore, for any $v \in V$ we have

$$\langle v, \underline{h} \rangle = \int_{[0, 1]^2} v(s, t) d(\tilde{h}_1'(s) \tilde{h}_2'(t)) \leq 0.$$

Consequently $\underline{h}$ belongs to the polar cone $\tilde{V}$ of V . In view of statement *c*) in Lemma 2.1 the proof follows if we show that g is orthogonal to $\underline{h}$. Since $\tilde{h}_i - h_i$ is orthogonal to $\tilde{h}_i, i = 1, 2$ (see Bischoff and Hashorva (2005)) we have

$$\begin{aligned}\langle g, \underline{h} \rangle &= \langle \tilde{h}_1 \times \tilde{h}_2 - h_1 \times h_2, \tilde{h}_1 \times \tilde{h}_2 \rangle \\ &= \langle \tilde{h}_1 \times (\tilde{h}_2 - h_2), \tilde{h}_1 \times \tilde{h}_2 \rangle - \langle (\tilde{h}_1 - h_1) \times h_2, \tilde{h}_1 \times \tilde{h}_2 \rangle \\ &= 0,\end{aligned}$$

hence the result follows. $\square$

Proof of Corollary 4.3: The proof follows easily by the assumptions on $u_i, i = 1, 2$. $\square$

6 Appendix

In this short section we provide two results for the Riemann-Stieltjes integral.

Let $f : [0, 1]^2 \rightarrow \mathbb{R}$ be a given function. If $f(s, t) = g(s, t) + g_1(s) + g_2(t)$ with $g \in BV_H([0, 1]^2)$ and g_1, g_2 two other functions, then h has bounded variation in the sense of Vitali (write $f \in BV_V([0, 1]^2)$). In fact f can be expressed as the difference of two real functions defined on $[0, 1]^2$ which generate a positive measure on $[0, 1]^2$. Thus the class of functions with bounded variation in the sense of Vitali consists of all real functions defined on $[0, 1]^2$ generating a finite signed measure.

If $g : [0, 1]^2 \rightarrow \mathbb{R}$ is continuous, then it is well-known that the Riemann-Stieltjes integral $\int_{[0, 1]^2} g(x, y) df(x, y)$ exists, provided that $f \in BV_V([0, 1]^2)$. In the next lemma we present an integration by parts formula, the case $f \in BV_H([0, 1]^2)$ is discussed in Lemma 1 in Móricz (2002).

Lemma 6.1 *Let $f, g : [0, 1]^2 \rightarrow \mathbb{R}$ be two given function. If g is continuous such that $g(s, t) = 0$ for all (s, t) in the boundary of $[0, 1]^2$ and $f \in BV_V([0, 1]^2)$, then the integration by parts formula for the Riemann-Stieltjes integral reads*

$$\int_{[0, 1]^2} g(x, y) df(x, y) = \int_{[0, 1]^2} f(x, y) dg(x, y). \quad (6.1)$$

Proof: The proof follows with similar arguments as in Lemma 2 in Móricz (2002), since the four single sums in the expression (3.8) therein are equal to 0 due to the fact that g vanishes on the boundary of $[0, 1]^2$. $\square$

Lemma 6.2 *Let $f, g : [0, 1]^2 \rightarrow \mathbb{R}$ be two given functions. Assume that g is absolute continuous with $g(s, t) = \int_{[0, s] \times [0, t]} h(x, y) d\lambda^2(dx, dy), s, t \in [0, 1]$. If $f \in BV_V([0, 1]^2)$ and f is almost surely continuous with respect to λ^2 , then we have*

$$\int_{[0, 1]^2} g(x, y) df(x, y) = \int_{[0, 1]^2} f(x, y) h(x, y) d\lambda^2(dx, dy). \quad (6.2)$$

Proof: The proof follows with similar arguments as in Lemma 3 in Móricz (2002). □

Acknowledgement: I would like to thank a Referee and Professor Wembo Li for several corrections and suggestions, Professor Móricz for sending [27], Professors Muhammad Aslam Noor and Wolfgang Bischoff for some insights on Hilbert spaces.

References

- [1] Adams, C.R., and Clarkson, J.A. (1934) Properties of functions $f(x, y)$ of bounded variation. *Trans. Amer. Math. Soc.* **36**, 711–730.
- [2] Bischoff, W., Miller, F., Hashorva, E., and Hüsler, J. (2003) Asymptotics of a boundary crossing probability of a Brownian bridge with general trend. *Meth. and Comp. Appl. Prob.* **5**, 3, 271–287.
- [3] Bischoff, W., Hashorva, E., Hüsler, J., and Miller, F. (2005) Analysis of a change-point regression problem in quality control by partial sums processes and Kolmogorov type tests. *Metrika* **62**, 85–98.
- [4] Bischoff, W., and Hashorva, E. (2005) A lower bound for boundary crossing probabilities of Brownian bridge/motion with trend *Stat. Prob. Lett.* **74**, 3, 265–271.
- [5] Bischoff, W., Hashorva, E., and Hüsler, J. (2007) An Asymptotic result for non-crossing probabilities of Brownian motion with trend. *Comm. Stat. Theory Methods*, **36**, 16, 2821–2828.
- [6] Borovkov, K., and Novikov, A. (2005) Explicit bounds for approximation rates of boundary crossing probabilities for the Wiener process. *J. Appl. Prob.* **42**, 82–92.
- [7] Csáki, E., Khoshnevisan, D., and Shi, Z. (2000) Boundary crossings and the distribution function of the maximum of Brownian sheet. *Stoch. Proc. Appl.* **90**, 1–18.
- [8] Fill, J., and Torcaso, F. (2004) Asymptotic analysis via Mellin transforms for small deviations in L_2 -norm of integrated Brownian sheets. *Probab. Theo. and Related Fields* **130**, 259–288.
- [9] Gao, F., Hannig, J., Lee, T.-Y., and Torcaso, F. (2004) Exact L_2 small balls of Gaussian processes. *J. Theoret. Probab.* **17**, 2, 503–520.
- [10] Gao, F., and Li, W.V. (2006) Logarithmic level comparison for small deviation probabilities. *J. Theoret. Probab.* **19**, 3, 535–556.
- [11] Gao, F., and Li, W.V. (2007) Small ball probabilities for the Slepian Gaussian fields. *Trans. Amer. Math. Soc.* **359**, 1339–1350.
- [12] Goovaerts, M.J., and Teunen, M. (1993) Boundary crossing results for the Brownian motion. *Blätter*, 197–205.
- [13] Hashorva, E. (2005a) Exact asymptotics for boundary crossing probabilities of Brownian motion with piecewise linear trend. *Elect. Comm. Prob.* **10**, 207–217.

- [14] Hashorva, E. (2005b) Asymptotics and bounds for multivariate Gaussian tails. *J. Theoret. Probab.* **18**, 1, 79–97.
- [15] Janssen, A., and Kunz, M. (2004) Boundary crossing probabilities for piece-wise linear boundary functions. *Comm. Stat. Theory Methods*, **33**, 7, 1445–1464.
- [16] Janssen, A., and Hülya, Ü. (2008) Regions of alternatives with high and low power for goodness-of-fit tests. *J. Stat. Plan. Inf.* **138**, 2526–2543.
- [17] Karol', A., Nazarov, A., Nikitin, Y. (2008) Small ball probabilities for Gaussian random fields and tensor products of compact operators. *Trans. Amer. Math. Soc.* **360**, 1443–1474.
- [18] Koning, A.J., and Protasov, V. (2003) Tail behaviour of Gaussian processes with applications to the Brownian pillow. *J. Mult. Anal.* **82**, 2, 370–397.
- [19] Khoshnevisan, D., and Pemantle, R. (2000) Sojourn times of Brownian sheet. *Periodica Math. Hung.* **41**, (1-2), 187–194
- [20] Kuelbs, J. (1976) A strong convergence theorem for Banach space valued random variables. *Ann. Probab.* **4**, 744–771.
- [21] Kuelbs, J., and Li, W.V. (1992) Small Ball Estimates for Brownian Motion and the Brownian Sheet. *J. Theoret. Probab.* **6**, 3, 547–577.
- [22] Kuelbs, J., and Li, W.V. (1994) Some large deviation results for gaussian measures. *Progr. Probab.* **35**, 251–270, Birkhäuser.
- [23] Ledoux, M. (1996) *Isoperimetry and Gaussian analysis. Lectures on probability theory and statistics. Lecture Notes in Math.* **1648**, Springer.
- [24] Li, W.V., and Kuelbs, J. (1998) Some shift inequalities for Gaussian measures. *Progr. Probab.* **43**, 233–243, Birkhäuser.
- [25] Li, W.V., and Shao, Q.M. (2001) Gaussian processes: Inequalities, small ball probabilities and applications. In *Stochastic Processes: Theory and Methods Handbook of Statistics* (C.R. Rao and D. Shanbhag, eds.) **19**, 533–597. North-Holland, Amsterdam.
- [26] Lifshits, M. A. (1995) *Gaussian random functions*. Mathematics and its Applications, 322. Kluwer Academic Publishers, Dordrecht.
- [27] Móricz, F. (2002) Order of magnitude of double Fourier coefficients of functions of bounded variation. *Analysis (Munich)* **22**, 335–345
- [28] Novikov, A.A., Frishling, V., and Kordzakhia, N. (1999) Approximations of boundary crossing probabilities for a Brownian motion. *J. Appl. Prob.* **36**, 1019–1030.
- [29] Pötzelberger, K., and Wang, L. (2001) Boundary crossing probability for Brownian motion. *J. Appl. Probab.* **38**, 152–164.

- [30] van der Vaart, A.W., and Wellner, J.A. (1996) *Weak convergence and empirical processes, with applications to statistics*. Springer, New York.
- [31] Wang, L., and Pötzelberger, K. (1997) Boundary crossing probability for Brownian motion and general boundaries. *J. Appl. Probab.* **34**, 54–65.